

ELECTROLYTE DISTURBANCES - HYPERKALAEMIA

Supporting information

This guideline has been prepared with reference to the following:

UK Kidney Association. Treatment of acute hyperkalaemia in adults. 2023. UK Kidney Association

<https://www.ukkidney.org/health-professionals/guidelines/treatment-acute-hyperkalaemia-adults-0>

Yartsev A. Response to 1L of Hartmann's compound sodium lactate (in Deranged Physiology). 2021.

<https://www.derangedphysiology.com/main/cicm-primary-exam/Chapter%20234/response-1l-hartmanns-compound-sodium-lactate>

NHS Improvement. Patient Safety Alert: Resources to support safe and timely management of hyperkalaemia. 2018

<https://www.england.nhs.uk/publication/patient-safety-alert-safe-and-timely-management-of-hyperkalaemia/>

Farkas, J. Myth-busting: Lactated Ringers is safe in hyperkalemia, and is superior to NS. 2014. EMCrit

<https://emcrit.org/pulmcrit/myth-busting-lactated-ringers-is-safe-in-hyperkalemia-and-is-superior-to-ns/>

If a patient on low molecular weight heparin develops hyperkalaemia, what should be done to discover whether this is drug-induced?

One method of identifying low molecular weight heparin (LMWH) as the cause of hyperkalemia is to reduce the dietary intake of potassium to see if the condition improves. The use of lactulose, chronological investigation and the stopping and starting of LMWH are other methods to discover if the condition is drug induced (Scalese, 2016).

Scalese MJ. Profound hyperkalemia associated with thromboprophylactic enoxaparin. Ther Adv Drug Saf. 2016;7: 120-1

Evidence Level: V

Last amended July 2024
Last reviewed January 2026

Not found an answer to your question? Wish to suggest an edit to this document?

Please contact the BCGP Clinical Effectiveness Librarian at bedsideclinicalguidelines@uuhm.nhs.uk