

URETHRAL CATHETERISATION

Supporting information

This guideline has been prepared with reference to the following:

NICE. Infection control, prevention of healthcare-associated infection in primary and community care. 2017. London. NICE

<http://www.nice.org.uk/guidance/cq139>

Bladder washouts do not prevent bacteriuria and are unnecessary as part of routine catheter care?

A 2017 systematic review of seven RCTs found that the evidence was not adequate to conclude if washouts were beneficial or harmful (Shepherd, 2017).

NICE Guidelines (2017) state that: "bladder instillations or washouts must not be used to prevent catheter associated infections".

A study of 120 routine catheter replacements not involving washouts (Bregenzer, 1997) found that bacteraemia occurred in only 5 cases (4.2%), which was not clinically significant.

Bladder washouts may be indicated in certain carefully-selected patients, either to remove blood clots from the bladder in a patient with haematuria or post TURP and TURBT, or to clear recurrent blockage, particularly if there is debris and/or stone formation. This should be performed only by a suitably experienced consultant (personal communication from Mr M Saxby, 16/12/06).

Bregenzer T, Frei R, Widmer AF, et al. Low risk of bacteraemia during catheter replacement in patients with long-term urinary catheters. *Arch Intern Med* 1997;157:521-5

Shepherd AJ, Mackay WG, Hagen S. Washout policies in long-term indwelling urinary catheterisation in adults.. Cochrane Database Syst Rev. 2017 Mar 6;3:CD004012

<https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD004012.pub5/full>

Evidence Level: I

Last amended March 2018
Last reviewed January 2026

Not found an answer to your question? Wish to suggest an edit to this document?

Please contact the BCGP Clinical Effectiveness Librarian at bedsideclinicalguidelines@uhnm.nhs.uk